

COINVARIANTS FOR LATTICE VOAS AND q -SUPERNOMIAL COEFFICIENTS

B. L. FEIGIN, S. A. LOKTEV, AND I. YU. TIPUNIN

ABSTRACT. We propose an alternative definition of q -supernomial coefficients as characters of coinvariants for one dimensional lattice vertex operator algebras. This gives a new formula for q -supernomial coefficients. Along the way we prove that the spaces of coinvariants form a bundle over the configuration space of complex points (the configuration space includes diagonals).

CONTENTS

1. Introduction	1	4. Degenerations of $\mathfrak{A}_p^{\text{ass}}$ and the main inequality	11
2. Lattice VOAs	3	5. Combinatorics of the characters	14
3. Coinvariants and Verlinde rules	7	6. Dimension and character formulas for the coinvariants	19
		References	21

1. INTRODUCTION

In this paper we study the spaces of conformal blocks for one dimensional lattice vertex operator algebras.

Spaces of conformal blocks or modular functor (see [S]) play an important role in conformal field theory. They can be defined for a wide class of vertex operator algebras and they parameterizes the states of the corresponding field theory. To avoid cumbersome notion let us describe our approach for affine algebras (which corresponds to the WZNW theory).

First, recall the definition of the space of conformal blocks. Let $\mathfrak{g}$ be a simple Lie algebra, by $\widehat{\mathfrak{g}}$ denote the central extension of the Lie algebra of $\mathfrak{g}$ -valued Laurent polynomials. Fix a level k , that is, identify the central element of $\widehat{\mathfrak{g}}$ with the positive integer k . As a geometric part of the data consider a complex curve C and a set of pairwise distinct points $p_1, \dots, p_n \in C$ with a local coordinate in a small vicinity of each point. Then define the Lie algebra $\mathfrak{g}^{C \setminus \{p_1, \dots, p_n\}}$ of $\mathfrak{g}$ -valued meromorphic functions on C with poles only at p_i . Clearly we have the inclusion

$$\mathfrak{g}^{C \setminus \{p_1, \dots, p_n\}} \hookrightarrow \widehat{\mathfrak{g}} \oplus \dots \oplus \widehat{\mathfrak{g}},$$

mapping any $\mathfrak{g}$ -valued function to its Laurent expansions at the points $p_1, \dots, p_n$. Another part of the data is a set of integrable representations $L_1, \dots, L_n$ of $\widehat{\mathfrak{g}}$ at level k . Their external tensor product is a representation of $\widehat{\mathfrak{g}} \oplus \dots \oplus \widehat{\mathfrak{g}}$ and, therefore, of $\mathfrak{g}^{C \setminus \{p_1, \dots, p_n\}}$. Then the space of conformal block is defined as the space of coinvariants

$$\langle L_1, \dots, L_n \rangle_{p_1, \dots, p_n}^C = (L_1 \boxtimes \dots \boxtimes L_n) / \mathfrak{g}^{C \setminus \{p_1, \dots, p_n\}},$$

i.e. the quotient of $L_1 \boxtimes \dots \boxtimes L_n$ by the action of $\mathfrak{g}^{C \setminus \{p_1, \dots, p_n\}}$. It is well known that these spaces are finite-dimensional and form a flat family over the module space of curves with punctures.

In [FFu] another construction is proposed. Let, as above, C be a curve, $p_1, \dots, p_n$ and p be pairwise distinct points on C . Then consider the subalgebra $\mathfrak{g}^{C \setminus p}(p_1, \dots, p_n) \subset \mathfrak{g}^{C \setminus p}$ of functions with zeroes at $p_1, \dots, p_n$ and the subalgebra $\mathfrak{g}_{\mathfrak{n}}^{C \setminus p}(p_1, \dots, p_n) \subset \mathfrak{g}^{C \setminus p}$ of functions whose values at $p_1, \dots, p_n$ belongs to the nilpotent Borel subalgebra $\mathfrak{n} \subset \mathfrak{g}$. Then for any integrable representation L at level k we have the natural isomorphisms (see [FKLMM1])

$$(1.1) \quad L/\mathfrak{g}_{\mathfrak{n}}^{C \setminus p}(p_1, \dots, p_n) \cong \bigoplus_{L_1, \dots, L_n} \langle L_1, \dots, L_n, L \rangle_{p_1, \dots, p_n, p}^C,$$

$$(1.2) \quad L/\mathfrak{g}^{C \setminus p}(p_1, \dots, p_n) \cong \bigoplus_{L_1, \dots, L_n} \pi(L_1) \otimes \dots \otimes \pi(L_n) \otimes \langle L_1, \dots, L_n, L \rangle_{p_1, \dots, p_n, p}^C,$$

where sum goes over all the sets of integrable representations at level k and $\pi(L_i)$ denotes the finite-dimensional representation of $\mathfrak{g}$ generated by the highest vector of L_i .

So the space of coinvariants of a single representation L contains the spaces of conformal blocks where L appears. An advantage of this construction is that these spaces of coinvariants is well defined if some of the points $p_1, \dots, p_n$ coincides. In this case we just consider zeroes with multiplicities. For multiple points we have not isomorphisms (1.1), (1.2), but it is easy to show that these spaces form a coherent sheaf over the variety $S^n C$ of sets of points on C .

First question is whether the dimension of the space of coinvariants preserves when some points coincides. Indeed, this question is local and it is enough to check this when C is the Riemann sphere. The answer is positive for $\mathfrak{g} = sl_2$ (see [FKLMM1] for $L/\mathfrak{g}^{C \setminus p}(p_1, \dots, p_n)$ and [FKLMM2] for $L/\mathfrak{g}_{\mathfrak{n}}^{C \setminus p}(p_1, \dots, p_n)$) and negative for $\mathfrak{g} = E_8$. Also there is the positive answer for the similar question about the $(2, m)$ -minimal models of the Virasoro algebras (see [FFr]). But the general case is far from been studied. Here we prove the equality of dimensions in the case of one dimensional lattice VOAs.

Another question is about the natural filtration on these spaces. Namely, the affine Lie algebra $\mathfrak{g}$ and its integrable representations are graded by the action of the energy element. But the subalgebras defined above are not homogeneous (except the case of Riemann sphere and $p_1 = \dots = p_n$), hence the spaces of coinvariants can be considered only as filtered spaces. Nevertheless we consider the Hilbert polynomials $\mathcal{H}(q)$ of the adjoint graded spaces and interpret them as a q -analog of the correspondent Verlinde numbers (which are dimensions of spaces of conformal blocks). This object is interesting even in the case of the Riemann sphere (and well studied only in this case). In [FL] there is a conjecture that expresses $\mathcal{H}(q)$ in terms of generalized q -supernomial coefficients. For $(2, m)$ -minimal models of the Virasoro algebras the polynomials $\mathcal{H}(q)$ take part in the Andrews-Gordon identities and therefore can be decomposed into sums of q -multinomial coefficients as well. In this paper we prove a formula (see Theorem 6.3, (ii)) which expresses $\mathcal{H}(q)$ for one-dimensional lattice VOAs as a sum of q -supernomial coefficients. Note that this formula can be considered as another definition of q -supernomials.

These two question are connected. Namely, suppose that the dimension preserves when the points coincides. Then in the case of the Riemann sphere it is enough to find $\mathcal{H}(q)$ for the

graded subalgebra with $p_1 = \dots = p_n$. It makes possible to degenerate the algebra together with the subalgebra into an abelian one or a rather simple VOA and obtain a fermionic type (in terms of [KKMM]) formula for $\mathcal{H}(q)$. Such a formula exists for $\mathfrak{g} = sl_2$ (see [FKLMM1], [FKLMM2], [FKLMM3]), for $(2, m)$ -minimal models of the Virasoro algebra (see [FFr]) and for one-dimensional lattice VOAs (see Theorem 6.3, (i)).

In more general cases (W -algebras, lattice VOAs, etc.) it is not evident which subalgebras should be chosen instead of $\mathfrak{g}^{C \setminus p}(p_1, \dots, p_n)$ and $\mathfrak{g}_n^{C \setminus p}(p_1, \dots, p_n)$. Let us explain what we are doing in the case of affine Lie algebras. Introduce some “elementary” graded subalgebras $\mathfrak{g}_i \subset \widehat{\mathfrak{g}}$. As the subalgebras are graded, they consist of $\mathfrak{g}$ -valued functions with some local conditions at the zero point. Then define the “fused” subalgebra $\mathfrak{g}_{i_1, \dots, i_n}^{C \setminus p}(p_1, \dots, p_n)$ of functions with pole only in p and local conditions in p_i prescribed by the subalgebras $\mathfrak{g}_i$. Then we have

$$(1.3) \quad L/\mathfrak{g}_{i_1, \dots, i_n}^{C \setminus p}(p_1, \dots, p_n) \cong \bigoplus_{L_1, \dots, L_n} (L_1/\mathfrak{g}_{i_1}) \otimes \dots \otimes (L_n/\mathfrak{g}_{i_n}) \otimes \langle L_1, \dots, L_n, L \rangle_{p_1, \dots, p_n, p}^C.$$

In particular, the subalgebra $\mathfrak{g}_n^{C \setminus p}(p_1, \dots, p_n)$ is fused from n copies of the affine nilpotent Borel subalgebra $\mathfrak{g}_n \subset \widehat{\mathfrak{g}}$ and the subalgebra $\mathfrak{g}^{C \setminus p}(p_1, \dots, p_n)$ is fused from n copies of the affine nilpotent parabolic subalgebra $\mathfrak{g}_0 \subset \widehat{\mathfrak{g}}$, so (1.1) and (1.2) follows from (1.3).

The paper is organized as follows.

In Sec. 2.1 we recall some facts about lattice vertex operator algebras (“VOA”) and their representations. In Sec. 2.2 we introduce an associative algebra that acts in each representation of the VOA. In this way representations of the VOA can be obtained as induced representations of the associative algebra.

In Sec. 3.1 we introduce spaces of coinvariants and give basic statements about their dimensions. In Sec. 3.2 elementary subalgebras are introduced and the fused subalgebras are studied.

Sec. 4 and 5 are rather technical. In Sec. 4 the character of some auxiliary space of coinvariants is calculated. In Sec. 5 we show that characters of these auxiliary spaces can be expressed in terms of q -supernomial coefficients. This allows us to prove the main results of the paper in Sec. 6.

We use the standard notation for the q -binomial coefficients

$$(1.4) \quad \begin{bmatrix} n \\ m \end{bmatrix}_q = \begin{cases} \frac{(q)_n}{(q)_{n-m}(q)_m}, & n \geq m \geq 0, \\ 0, & \text{otherwise,} \end{cases} \quad \text{where } (q)_n = (1-q) \dots (1-q^n).$$

2. LATTICE VOAS

2.1. Recall that a lattice Vertex Operator Algebra (VOA) is determined by the following data: $\mathbb{R}^n$, $\langle \cdot, \cdot \rangle$, Γ , where $\langle \cdot, \cdot \rangle$ is a scalar product in $\mathbb{R}^n$, Γ is a lattice in $\mathbb{R}^n$ and $\langle \cdot, \cdot \rangle$ takes integer values on vectors from Γ . Such an algebra is generated by vertex operators $V(\mathbf{x}, z)$, $\mathbf{x} \in \Gamma$.

In this paper we concentrate at one dimensional lattice VOAs with positively defined scalar product. Let α be the basis vector of Γ and $\langle \alpha, \alpha \rangle = p$, $p \in \mathbb{N}$. Denote this VOA by $\mathfrak{A}_p$.

for $p = 1, 2, 3, 4, \dots$. The algebra $\mathfrak{A}_p$ is generated by $\mathcal{G}^+(z) = V(\alpha, z)$ and $\mathcal{G}^-(z) = V(-\alpha, z)$. The generators $\mathcal{G}^\pm$ are bosonic ($\mathcal{G}^\pm(z)\mathcal{G}^\pm(w) - \mathcal{G}^\pm(w)\mathcal{G}^\pm(z) = 0$) whenever p is even and fermionic ($\mathcal{G}^\pm(z)\mathcal{G}^\pm(w) + \mathcal{G}^\pm(w)\mathcal{G}^\pm(z) = 0$) otherwise and satisfy the following relations

$$(2.1) \quad (\mathcal{G}^+(z))^2 = 0, \quad (\partial\mathcal{G}^+(z))^2 = 0, \quad \dots, \quad (\partial^{p-3}\mathcal{G}^+(z))^2 = 0,$$

$$(2.2) \quad (\mathcal{G}^-(z))^2 = 0, \quad (\partial\mathcal{G}^-(z))^2 = 0, \quad \dots, \quad (\partial^{p-3}\mathcal{G}^-(z))^2 = 0,$$

whenever p is even and

$$(2.3) \quad \partial\mathcal{G}^+(z)\mathcal{G}^+(z) = 0, \quad \partial^3\mathcal{G}^+(z)\mathcal{G}^+(z) = 0, \quad \dots, \quad \partial^{p-2}\mathcal{G}^+(z)\mathcal{G}^+(z) = 0,$$

$$(2.4) \quad \partial\mathcal{G}^-(z)\mathcal{G}^-(z) = 0, \quad \partial^3\mathcal{G}^-(z)\mathcal{G}^-(z) = 0, \quad \dots, \quad \partial^{p-2}\mathcal{G}^-(z)\mathcal{G}^-(z) = 0,$$

whenever p is odd.

Introduce currents $\mathcal{H}^m$ for $0 \leq m \leq p-2$ as the currents appear in the OPE

$$(2.5) \quad \mathcal{G}^+(z)\mathcal{G}^-(w) = \frac{\frac{1}{p}}{(z-w)^p} + \frac{\mathcal{H}^0}{(z-w)^{p-1}} + \frac{\mathcal{H}^1}{(z-w)^{p-2}} + \dots + \frac{\mathcal{H}^{p-2}}{z-w} + \text{regular terms}.$$

Currents $\mathcal{H}^0$ and $\mathcal{H}^1$ correspond to $u(1)$ -current and energy-momentum tensor respectively. Choose Virasoro $\mathcal{T}$ so that $\mathcal{G}^+$ and $\mathcal{G}^-$ be primary fields with conformal weights $p-1$ and 1 respectively

$$(2.6) \quad \mathcal{T} = \mathcal{H}^1 - \frac{p-1}{2}\partial\mathcal{H}^0.$$

The mode decomposition corresponding the choice (2.6) is

$$(2.7) \quad \mathcal{G}^+(z) = \sum_{n \in \mathbb{Z}} \mathcal{G}_n^+ z^{-n-p+1}, \quad \mathcal{G}^-(z) = \sum_{n \in \mathbb{Z}} \mathcal{G}_n^- z^{-n-1},$$

$$(2.8) \quad \mathcal{H}^m(z) = \sum_{n \in \mathbb{Z}} \mathcal{H}_n^m z^{-n-m-1}, \quad \mathcal{T}(z) = \sum_{n \in \mathbb{Z}} \mathcal{L}_n z^{-n-2}.$$

This gives the following commutation relations

$$(2.9) \quad [\mathcal{L}_n, \mathcal{G}_i^+] = ((p-2)n - i)\mathcal{G}_{i+n}^+, \quad [\mathcal{L}_n, \mathcal{G}_i^-] = -i\mathcal{G}_{i+n}^-, \quad [\mathcal{L}_0, \mathcal{H}_i^m] = -i\mathcal{H}_i^m.$$

The algebra $\mathfrak{A}_p$ admits the following family of automorphisms (spectral flow) U_θ :

$$(2.10) \quad \begin{aligned} \mathcal{G}_n^+ &\mapsto \mathcal{G}_{n+\theta}^+, & \mathcal{G}_n^- &\mapsto \mathcal{G}_{n-\theta}^-, \\ \mathcal{H}_n^0 &\mapsto \mathcal{H}_n^0 + \theta \frac{1}{p} \delta_{n,0}, & \mathcal{H}_n^1 &\mapsto \mathcal{H}_n^1 + \theta \mathcal{H}_n^0 + \frac{1}{2} \theta(\theta-1) \frac{1}{p} \delta_{n,0}, \\ \mathcal{H}_n^m &\mapsto \mathcal{H}_n^m + \theta \mathcal{H}_n^{m-1} + \frac{1}{2} \theta(\theta-1) \mathcal{H}_n^{m-2} + \frac{1}{6} \theta(\theta-1)(\theta-2) \mathcal{H}_n^{m-3} + \dots \\ & & & + \frac{1}{m!} \theta(\theta-1)(\theta-2) \dots (\theta-m) \frac{1}{p} \delta_{n,0}, \end{aligned}$$

where $\theta \in \mathbb{Z}$. The automorphisms with $\theta \in p\mathbb{Z}$ play the role of the affine part of the Weyl group.

Note that in the case $p = 1$ the algebra $\mathfrak{A}_p$ can be considered as the infinite-dimensional Clifford algebra, in the case $p = 2$ as $\widehat{sl}_2$ at level $k = 1$ and in the case $p = 3$ as the $N=2$ -superconformal algebra with the central charge $c = \frac{1}{3}$.

Recall some facts about representations of lattice VOAs. The category of representations of any lattice VOA is semisimple, that is, any representation decomposes into the direct sum of irreducible ones. The irreducible representations are enumerated by the quotient $\Gamma^\vee/\Gamma$, where $\Gamma^\vee$ is the dual lattice. Namely, the representation corresponding to the element $\mathbf{y} \in \Gamma^\vee$ is the space of descendants of $V(\mathbf{y}, 0)$, and if two elements of $\Gamma^\vee$ differs by an element of Γ then the corresponding representations are isomorphic. It is also known that these representations form a minimal model with the Verlinde algebra isomorphic to the group ring of $\Gamma^\vee/\Gamma$.

Let us describe it in detail for the one-dimensional case. Denote the representation corresponding to $V(-\frac{r}{\alpha}, 0)$ by $\mathbf{l}_r$ with $0 \leq r \leq p-1$. This representation contains the highest weight vector $|p; r\rangle$ satisfying the following conditions

$$(2.11) \quad \mathcal{G}^+_i |p; r\rangle = \mathcal{G}^-_j |p; r\rangle = 0, \quad i \geq -p + r + 2, \quad j \geq -r, \\ \mathcal{H}^0_0 |p; r\rangle = -\frac{n}{p} |p; r\rangle, \quad \mathcal{L}_0 |p; r\rangle = \frac{n}{2p} (r - p + 2) |p; r\rangle.$$

Another description of $\mathbf{l}_r$ is as follows

$$(2.12) \quad \mathbf{l}_r = \bigoplus_{m \in \mathbb{Z}} \mathbf{l}_r(m)$$

where $\mathbf{l}_r(m)$ is the Fock module over $\mathcal{H}^0$ with the highest weight vector $|p; r + pm\rangle$. In this realization $\mathcal{G}^\pm : \mathbf{l}_r(m) \rightarrow \mathbf{l}_r(m \pm 1)$ act as vertex operators. We call the representation $\mathbf{l}_0$ a *vacuum* representation.

Representations $\mathbf{l}_r$ with $0 \leq r \leq p-1$ compose a minimal model and the Verlinde algebra (the fusion algebra) is

$$(2.13) \quad \mathbf{l}_r \cdot \mathbf{l}_s = \mathbf{l}_{r+s \bmod p}$$

The vacuum representation $\mathbf{l}_0$ is the unit in the Verlinde algebra. Note also that the module $\mathbf{l}_r$ is isomorphic to the contragradient to $\mathbf{l}_{p-r}$. This equips the Verlinde algebra with the scalar product.

Automorphisms $\mathbf{U}_\theta$ naturally act on the set of representations $\mathbf{l}_r$ as

$$(2.14) \quad \mathbf{U}_\theta \mathbf{l}_r = \mathbf{l}_{r+\theta \bmod p}.$$

In particular, the action of $\mathbf{U}_1$ on the set of representations $\mathbf{l}_r$ coincides with the fusion with $\mathbf{l}_1$.

Each space $\mathbf{l}_r$ is bi-graded by operators $\mathcal{H}^0_0$ and $\mathcal{L}_0$. Characters of representations $\mathbf{l}_r$ corresponding to this bi-grading are by definition

$$(2.15) \quad \text{ch } \mathbf{l}_r(q, z) = \text{tr } z^{\mathcal{H}^0_0} q^{\mathcal{L}_0},$$

where the trace is calculated over $\mathbf{l}_r$ and (2.12) implies

$$(2.16) \quad \text{ch } \mathbf{l}_r(q, z) = \frac{z^{-\frac{r}{p}} q^{\frac{r}{2p}(r-p+2)}}{(q)_\infty} \sum_{n \in \mathbb{Z}} z^n q^{\frac{p}{2}(n^2+n)-n(r+1)}$$

For any graded subspace an analogous series can be considered. We will denote them by the symbol ch in front of the symbol of the subspace.

2.2. Representations of $\mathfrak{A}_p$ can be considered from a more algebraic viewpoint. Operators $\mathcal{G}^\pm_n$ act on each representation $\mathfrak{l}_r$ therefore the associative algebra generated by $\mathcal{G}^\pm_n$ acts on each $\mathfrak{l}_r$. Denote this algebra by $\mathfrak{A}_p^{\text{ass}}$ and let us describe its defining relations.

Note first of all that $\mathfrak{A}_p$ contains a Virasoro algebra hence the algebra of vector fields on the circle acts by derivatives (commutators with $\mathcal{L}_j$) on $\mathfrak{A}_p^{\text{ass}}$. By $\mathcal{F}_\lambda$ denote the representation of the algebra of vector fields in weight λ tensor fields $\mathbb{C}[z, z^{-1}](dz)^\lambda$. Then the space $\{\mathcal{G}^+_i, i \in \mathbb{Z}\} = W_+$ is isomorphic to $\mathcal{F}_{2-p}$ and $\{\mathcal{G}^-_i, i \in \mathbb{Z}\} = W_-$ is isomorphic to $\mathcal{F}_0$. This isomorphism is given by the identification $\mathcal{G}^+_i = z_1^{i-p+2}(dz_1)^{2-p}$, $\mathcal{G}^-_i = z_2^i(dz_2)^0$.

Generators $\mathcal{G}^\pm_i$ commute whenever p is even, anticommute otherwise and in both cases satisfy quadratic relations

$$(2.17) \quad \sum_{\alpha+\beta=n} p(\alpha, \beta) \mathcal{G}^\pm_\alpha \mathcal{G}^\pm_\beta = 0, \quad n \in \mathbb{Z},$$

for any polynomial $p(\alpha, \beta)$ of degree not bigger than $p-2$.

The relations between $\mathcal{G}^+_i$ and $\mathcal{G}^-_j$ are also quadratic. Introduce the space W_0 of all commutators of $\mathcal{G}^+_i$ and $\mathcal{G}^-_j$ for even p and anticommutators for odd p . Let us describe W_0 as a quotient of the tensor product $W_+ \otimes W_-$.

We identify $W_+ \otimes W_-$ with the space of Laurent polynomials $\mathbb{C}[z_1, z_1^{-1}, z_2, z_2^{-1}](dz_1)^{2-p}(dz_2)^0$. Then introduce the subspaces

$$W_0^{(j)} = \mathbb{C}[z_1, z_1^{-1}, z_2, z_2^{-1}](z_1 - z_2)^{p-2-j}(dz_1)^{2-p}(dz_2)^0, \quad j \in \mathbb{Z}.$$

Note that the quotient $W_0^{(j)}/W_0^{(j-1)}$ is isomorphic to $\mathcal{F}_{-j}$ as a module over the algebra of vector fields.

The OPE (2.5) implies that the space $W_0^{(-2)}$ acts on any $\mathfrak{l}_r$ by zero and $W_0^{(-1)}$ acts by a scalar. The scalar can be found as follows. We know that $W_0^{(-1)}/W_0^{(-2)} \simeq \mathcal{F}_1$, but $\mathcal{F}_1$ is the space of 1-forms, so there exists the functional

$$(2.18) \quad \beta : W_0^{(-1)} \longrightarrow \mathcal{F}_1 \xrightarrow{\text{res}} \mathbb{C}$$

Let $K \subset W_0^{(-1)}$ be the kernel of the mapping β . Then K is a subspace in $W_+ \otimes W_-$ and we have $W_0 = (W_+ \otimes W_-)/K$.

Theorem 2.1. *The defining relations in $\mathfrak{A}_p^{\text{ass}}$ are (2.17) and relations between W_+ and W_- following from (2.18).*

We don't use this Theorem anywhere in the paper, so let us omit the proof. But this consideration gives us the following description of $\mathcal{H}^i_s$ in terms of Laurent polynomials

$$(2.19) \quad \mathcal{H}^i_s = z_1^{-p+2} z_2^{s+i} (z_1 - z_2)^{p-i-2} (dz_1)^{p-2} (dz_2)^0$$

This notion is very convenient for us. In particular, automorphism (2.10) acts on the space of Laurent polynomials as the multiplication with $(z_1/z_2)^\theta$.

One can show that the category of representations of $\mathfrak{A}_p^{\text{ass}}$ with highest weight is equivalent to the category of representations of $\mathfrak{A}_p$. So irreducible representations of $\mathfrak{A}_p^{\text{ass}}$ with highest weight are precisely $\mathfrak{l}_r$.

Define for the algebra $\mathfrak{A}_p^{\text{ass}}$ some class of representations. Each of these representations contains a cyclic vector that is annihilated by some subspaces in $W_+ \oplus W_0 \oplus W_-$, which we now describe. Let $\mathbf{N} = (N_+, N_-; N_0, N_1, \dots, N_{p-2})$ be a $p+1$ -dimensional vector with integer components that satisfy

$$(2.20) \quad 0 \leq N_0 \leq N_1 \leq \dots \leq N_{p-2} \leq N_+ + N_-.$$

We define the subspace $W(\mathbf{N}) = W_+(\mathbf{N}) \oplus W_0(\mathbf{N}) \oplus W_-(\mathbf{N})$ as follows

$$\begin{aligned} W_+(\mathbf{N}) &= \mathbb{C}[z_1]z_1^{N_+}(dz_1)^{2-p} \subset \mathbb{C}[z_1, z_1^{-1}](dz_1)^{2-p}, \\ W_-(\mathbf{N}) &= \mathbb{C}[z_2]z_2^{N_-}(dz_2)^0 \subset \mathbb{C}[z_2, z_2^{-1}](dz_2)^0 \end{aligned}$$

and the space $W_0(\mathbf{N})$ is the sum (not direct!) of the images of the subspaces

$$W_0^{(j)}(\mathbf{N}) = \mathbb{C}[z_1, z_2]z_1^{N_j - N_-}z_2^{N_-}(z_1 - z_2)^{p-j-2}(dz_1)^{2-p}(dz_2)^0$$

for $j = 0 \dots p-2$ under the mapping $W_+ \otimes W_- \rightarrow W_0$.

Note that if $N_0 = N_1 = \dots = N_{p-2} = N_+ + N_-$ then $W_0(\mathbf{N})$ is the subspace of (anti)commutators of $W_+(\mathbf{N})$ and $W_-(\mathbf{N})$.

Clearly we have $U_\theta(W(\mathbf{N})) = W(\mathbf{N} + \theta \mathbf{u})$, where

$$(2.21) \quad \mathbf{u} = (1, -1; 0, \dots, 0)$$

Proposition 2.2. *The irreducible representation $\mathfrak{l}_r$ is induced from the subspace $W(\mathbf{ru})$.*

Example 2.3. Let $p = 2$. Then we can identify $\mathcal{G}_i^+ = e \cdot t^i$, $\mathcal{G}_i^- = f \cdot t^i$, $\mathcal{H}_i^0 = h \cdot t^i$, where e, h, f is the standard basis of sl_2 . So our VOA is isomorphic to $\widehat{sl_2}$ at level 1. Then we have

$$(2.22) \quad W((N_+, N_-, N_0)) = e \cdot t^{N_+} \mathbb{C}[t] \oplus h \cdot t^{N_0} \mathbb{C}[t] \oplus f \cdot t^{N_-} \mathbb{C}[t]$$

So for $k = 1$ we cover the cases considered in [FKLMM1, FKLMM2, FKLMM3].

Let $R(\mathbf{N})$ be the representation induced from the trivial representation of the subalgebra generated by the subspace $W(\mathbf{N})$. We know that the representation $R(\mathbf{N})$ is a direct sum of $\mathfrak{l}_r$. The multiplicity of $\mathfrak{l}_r$ in $R(\mathbf{N})$ is the dimension of the space $\text{Hom}_{\mathfrak{A}_p^{\text{ass}}}(R(\mathbf{N}), \mathfrak{l}_r)$, which is isomorphic to the space of $W(\mathbf{N})$ -invariants in $\mathfrak{l}_r$.

3. COINVARIANTS AND VERLINDE RULES

3.1. It is more convenient to study the dual space to the space of $W(\mathbf{N})$ -invariants in $\mathfrak{l}_r$ namely the space of $W(\mathbf{N})$ -coinvariants in $\mathfrak{l}_r^*$, which is by definition $\mathfrak{l}_r^*/W(\mathbf{N})\mathfrak{l}_r^*$. Note that the dual representation $\mathfrak{l}_r^*$ differs from $\mathfrak{l}_r$ only by exchanging positive and negatives modes.

The space $W(\mathbf{N})$ can be deformed, i.e. can be included in a family of subspaces of W depending on a set of parameters that are points on $\mathbb{C}$. Given k vectors $\mathbf{N}^i = (N_+^i, N_-^i; N_0^i, N_1^i, \dots, N_{p-2}^i)$ satisfying (2.20) and given a vector $\mathbf{a} = (a^1, \dots, a^k) \in \mathbb{C}^k$ we define $W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ as the direct sum of the spaces $W_{\pm}(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ and $W_0(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$, where

$$W_+(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a}) = \mathbb{C}[z_1](dz_1)^{2-p} \prod_{i=1}^k (z_1 - a_i)^{N_+^i} \subset \mathbb{C}[z_1, z_1^{-1}](dz_1)^{2-p},$$

$$W_-(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a}) = \mathbb{C}[z_2](dz_2)^0 \prod_{i=1}^k (z_2 - a_i)^{N_-^i} \subset \mathbb{C}[z_2, z_2^{-1}](dz_2)^0$$

and the space $W_0(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ is the sum (not direct!) of the images of the subspaces

$$W_0^{(j)}(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a}) = \mathbb{C}[z_1, z_2](z_1 - z_2)^{p-j-2}(dz_1)^{2-p}(dz_2)^0 \prod_{i=1}^k (z_1 - a_i)^{N_+^i - N_-^i} (z_2 - a_i)^{N_-^i}$$

for $j = 0 \dots p-2$ under the mapping $W_+ \otimes W_- \rightarrow W_0$.

Then $W(\mathbf{N}) = W(\mathbf{N}^1, \mathbf{N}^2, \dots, \mathbf{N}^k; 0)$ for $\mathbf{N} = \mathbf{N}^1 + \dots + \mathbf{N}^k$, so the family of subalgebras $W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ is a deformation of $W(\mathbf{N})$.

Such a deformation can be applied to the calculation of dimensions for the coinvariants. To simplify the notation denote by $\mathbf{l}_r[W]$ the space of coinvariants $\mathbf{l}_r^*/W\mathbf{l}_r^*$.

Theorem 3.1. *Suppose that points $a^1, a^2, \dots, a^k$ are pairwise distinct. Then the dimension of $\mathbf{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})]$ coincides with the coefficient before $\mathbf{l}_r$ in the following expression*

$$(3.1) \quad \prod_{i=1}^k (\dim \mathbf{l}_0[W(\mathbf{N}^i)] \cdot \mathbf{l}_0 + \dots + \dim \mathbf{l}_{p-1}[W(\mathbf{N}^i)] \cdot \mathbf{l}_{p-1}),$$

where product means the Verlinde multiplication.

Proof. First recall the fusion procedure in the case of one input and k outputs.

Suppose that $\mathbf{N}^1, \dots, \mathbf{N}^k$ and $\mathbf{M}^1, \dots, \mathbf{M}^k$ are two sets of vectors such that $\mathbf{N}_j^i \geq \mathbf{M}_j^i$ for any $i = 1 \dots k, j = +, -, 0, \dots, p-2$. Then we have the natural projection $\mathbf{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})] \rightarrow \mathbf{l}_r[W(\mathbf{M}^1, \dots, \mathbf{M}^k; \mathbf{a})]$. Therefore the dual spaces form an inductive system and we can consider the limit

$$\mathbf{l}_r[\mathbf{a}] = \varinjlim (\mathbf{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})])^*.$$

The action of the algebra $\mathfrak{A}_p^{\text{ass}}$ on $\mathbf{l}_r$ can be extended to the action of the algebra $\mathfrak{A}_p^{\text{ass}} \oplus \dots \oplus \mathfrak{A}_p^{\text{ass}}$ on the limit $\mathbf{l}_r[\mathbf{a}]$, where summands correspond to points $a^1, \dots, a^k$.

The statement that the set of representations $\mathbf{l}_r$ form a minimal model means that the representation $\mathbf{l}_r[\mathbf{a}]$ decomposes into the direct sum of external tensor products

$$\mathbf{l}_r[\mathbf{a}] = \bigoplus_{\substack{0 \leq s_1, \dots, s_k < p \\ \mathbf{l}_{s_1} \cdot \mathbf{l}_{s_2} \cdot \dots \cdot \mathbf{l}_{s_k} = \mathbf{l}_r}} \mathbf{l}_{s_1} \boxtimes \dots \boxtimes \mathbf{l}_{s_k}$$

Emphasize that the sum goes over the set of representations whose product in the Verlinde algebra is $\mathbf{l}_r$.

Clearly, the space $\mathfrak{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})]$ is dual to the space of $W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ -invariants in $\mathfrak{l}_r[\mathbf{a}]$. As the subspace $W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})$ is dense with respect to the topology of direct limit in the subspace $W(\mathbf{N}^1) \oplus \dots \oplus W(\mathbf{N}^k) \subset \mathfrak{A}_p^{\text{ass}} \oplus \dots \oplus \mathfrak{A}_p^{\text{ass}}$, we have

$$\dim \mathfrak{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})] = \sum_{\substack{0 \leq s_1, \dots, s_k < p \\ \mathfrak{l}_{s_1} \cdot \mathfrak{l}_{s_2} \cdots \mathfrak{l}_{s_k} = \mathfrak{l}_r}} \dim \mathfrak{l}_{s_1}[W(\mathbf{N}^1)] \cdots \dim \mathfrak{l}_{s_k}[W(\mathbf{N}^k)],$$

that is equivalent to the statement of the theorem. $\square$

Here we have a family of subalgebras parametrized by complex numbers $a^1, \dots, a^k$. The deformation argument (see for example [FKLMM2]) implies that dimension of the space of coinvariants with respect to a special subalgebra is not less than one with respect to a generic subalgebra.

Proposition 3.2. *The dimension of $\mathfrak{l}_r[W(\mathbf{N}^1, \dots, \mathbf{N}^k; \mathbf{a})]$ is minimal whenever a^i are pairwise distinct and maximal whenever a^i are equal each to other.*

3.2. Introduce elementary spaces $W_{i,j} = W(\mathbf{N}^{i,j})$ with

$$\mathbf{N}^{i,j} = (i - j, j; 1, 2, \dots, i - 1, i, \dots, i)$$

and $0 \leq i \leq p, j \in \mathbb{Z}$.

Proposition 3.3. *We have*

$$(3.2) \quad \dim \mathfrak{l}_r[W_{i,j}] = \#\{n | 0 \leq pn + j + r \leq i\}$$

In other words,

$$\dim \mathfrak{l}_r[W_{i,j}] = \begin{cases} 2 & i = p, (r + j) \bmod p = 0 \\ 1 & \text{otherwise} \\ 0 & (r + j) \bmod p > i \end{cases}$$

Proof. Note that $W_{i,j}$ contains the subalgebra $\mathcal{H}_{\geq 1}^0$ therefore the space of $W_{i,j}$ -invariants in $\mathfrak{l}_r$ is spanned by several vectors $|p; r + pn\rangle, n \in \mathbb{Z}$. Taking the invariants of $\mathcal{G}_{\geq i-j-p+2}^+$ and $\mathcal{G}_{\geq j}^-$ selects the vectors $|p; r + pn\rangle$ with $-j \leq r + pn \leq i - j$. We need to show that these vectors are invariant.

Note that if $-j \leq r + pn \leq i - j$ then $W_{i,j} \subset W_{0, -(r+pn)}$. And we know that the vector $|p; s\rangle$ is annihilated by the subspace $W_{0, -s}$. $\square$

According to (3.1), (3.2) and (2.13) we define the positive integers $d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k]$ by the following formula

$$(3.3) \quad \prod_{s=1}^k (\xi^{-j^s} + \xi^{-j^s+1} + \dots + \xi^{-j^s+i^s}) = \sum_{r=0}^{p-1} d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k] \cdot \xi^r,$$

where ξ is a primitive root of unity of degree p .

Then by Theorem 3.1 and Proposition 3.2 we have

$$(3.4) \quad \dim \mathfrak{l}_r[W(\mathbf{N}^{i^1, j^1}, \dots, \mathbf{N}^{i^k, j^k}; 0)] \geq \dim \mathfrak{l}_r[W(\mathbf{N}^{i^1, j^1}, \dots, \mathbf{N}^{i^k, j^k}; \mathbf{a})] \geq d_{p,r}[i^1, j^1; \dots; i^k, j^k]$$

for any $\mathbf{a} \in \mathbb{C}^k$.

One aim of this paper is to show that (3.4) is indeed an equality for arbitrary vector $\mathbf{a} \in \mathbb{C}^k$ (see Theorem 6.1).

It is easy to see from (3.3) that $d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k]$ is a sum with the step p of supernomial coefficients. To write this sum introduce the $m \times m$ matrix¹

$$(3.5) \quad T_m = \begin{pmatrix} 2 & -1 & 0 & 0 & \dots & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & \dots & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & \dots & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 & -1 & 1 \end{pmatrix}$$

so $(T_m^{-1})_{ij} = \min\{i, j\}$ for $1 \leq i, j \leq m$.

With the vector $\mathbf{N} = (N_+, N_-; N_0, N_1, \dots, N_{p-2})$ we associate the other vector

$$\mathbf{N}' = (N_0, N_1, \dots, N_{p-2}, N_+ + N_-).$$

Define the vector $\mathbf{L} = (L_1, L_2, \dots, L_p)$ as follows

$$(3.6) \quad \mathbf{L} = \mathbf{N}' T_p.$$

Proposition 3.4. *Let $\mathbf{N} = \sum_{s=1}^k \mathbf{N}^{i^s, j^s}$ and $\mathbf{L} = \mathbf{N}' T_p$. Then*

$$(3.7) \quad d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k] = \sum_{a \in \mathbb{Z}} \binom{\mathbf{L}}{pa + N_- - r}^{p+1},$$

In the case where at least one of $i^s = p - 1$ sum (3.7) can be easily calculated.

Remark 3.5. We have

$$(3.8) \quad d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k] = 2^{L_1} 3^{L_2} \dots (p-1)^{L_{p-2}} p^{L_{p-1}-1} (p+1)^{L_p}$$

whenever $L_{p-1} \geq 1$ or equivalently at least one i^s is equal to $p - 1$.

At last let us describe the class of subalgebras $W(\mathbf{N})$ which can be deformed in this way.

Proposition 3.6. *Suppose that all the components of the vector $\mathbf{L} = \mathbf{N}' T_{p-1}$ are non-negative. Then we have*

$$W(\mathbf{N}) \simeq W(\mathbf{N}^{i^1, j^1}, \mathbf{N}^{i^2, j^2}, \dots, \mathbf{N}^{i^k, j^k}; 0)$$

for some $i^1, \dots, i^k; j^1, \dots, j^k$.

¹See [SW]

4. DEGENERATIONS OF $\mathfrak{A}_p^{\text{ass}}$ AND THE MAIN INEQUALITY

4.1. Let I be a set with ℓ elements. Let $\mathbf{e}_a$ with $a \in I$ be the standard basis in $\mathbb{Z}^\ell$. We write elements $\mathbf{x}$ from $\mathbb{Z}^\ell$ as rows. Let $\mathbf{x} \cdot \mathbf{y}$ denote the standard scalar product in $\mathbb{Z}^\ell$.

Let $\Gamma(A)$ be the lattice with the basis $\mathbf{e}_a$ and the scalar product determined by a symmetric matrix A

$$(4.1) \quad \langle \mathbf{e}_a, \mathbf{e}_b \rangle = A_{ab} \in \mathbb{Z}, \quad a, b \in I.$$

Consider the algebra $\overline{\mathfrak{B}}(A)$ generated by vertex operators $V(\mathbf{e}_a, z)$, $a \in I$. Emphasize that we do not involve vertex operators corresponding to $-\mathbf{e}_a$. Let us fix the decomposition of $V(\mathbf{e}_a, z)$ into formal power series

$$(4.2) \quad V(\mathbf{e}_a, z) = \sum_{n \in \mathbb{Z}} V(\mathbf{e}_a)_n z^{-n-1}.$$

The algebra $\overline{\mathfrak{B}}(A)$ is naturally $\Gamma(A)$ -graded. Decomposition (4.2) determines another one grading by operator $\mathcal{L}_0$

$$(4.3) \quad [\mathcal{L}_0, V(\mathbf{e}_a)_n] = -nV(\mathbf{e}_a)_n.$$

This algebra acts in the space

$$(4.4) \quad \mathbf{H} = \bigoplus_{\mathbf{y} \in \Gamma^\vee(A)} \mathbf{H}(\mathbf{y}),$$

where $\mathbf{H}(\mathbf{y})$ is the Fock module with the momentum $\mathbf{y}$ and $\Gamma^\vee(A)$ is the lattice dual to $\Gamma(A)$. The representation $\mathbf{H}$ contains the subrepresentation $\mathbf{J}_{\mathbf{y}}$ generated from the highest weight vector $|\mathbf{y}\rangle$ in $\mathbf{H}(\mathbf{y})$. The representation $\mathbf{J}_{\mathbf{y}}$ plays a role of Verma module, namely, it is maximal among representation with the same annihilation conditions satisfied by the vector $|\mathbf{y}\rangle$. Clearly $\mathbf{J}_{\mathbf{y}}$ admits the $\Gamma^\vee(A)$ -grading; to fix it let us set $\deg |\mathbf{y}\rangle = 0$.

The dual space $\mathbf{J}_{\mathbf{y}}^*$ can be identified with the space of correlation functions. Namely, any homogeneous linear functional $\langle \theta | \in \mathbf{J}_{\mathbf{y}}^*$ of degree $\sum n_i \mathbf{e}_i$ determines and is determined by the function

$$(4.5) \quad Y_{\theta, \mathbf{y}}(x_1^1, x_2^1, \dots, x_{n_1}^1; x_1^2, x_2^2, \dots, x_{n_2}^2; \dots; x_1^\ell, x_2^\ell, \dots, x_{n_\ell}^\ell) = \\ \langle \theta | V(\mathbf{e}_1, x_1^1) V(\mathbf{e}_1, x_2^1) \dots V(\mathbf{e}_1, x_{n_1}^1) V(\mathbf{e}_2, x_1^2) V(\mathbf{e}_2, x_2^2) \dots V(\mathbf{e}_2, x_{n_2}^2) \dots \\ V(\mathbf{e}_\ell, x_1^\ell) V(\mathbf{e}_\ell, x_2^\ell) \dots V(\mathbf{e}_\ell, x_{n_\ell}^\ell) |\mathbf{y}\rangle.$$

Proposition 4.1. *Correlation functions $Y_{\theta, \mathbf{y}}$ are products of the polynomial*

$$(4.6) \quad (x_1^1 x_2^1 \dots x_{n_1}^1)^{w_1} (x_1^2 x_2^2 \dots x_{n_2}^2)^{w_2} \dots (x_1^\ell x_2^\ell \dots x_{n_\ell}^\ell)^{w_\ell} \prod_{\substack{1 \leq a \leq \ell \\ 1 \leq i < j \leq n_a}} (x_i^a - x_j^a)^{A_{aa}} \prod_{\substack{1 \leq a < b \leq \ell \\ 1 \leq i \leq n_a \\ 1 \leq j \leq n_b}} (x_i^a - x_j^b)^{A_{ab}}$$

and a polynomial symmetric in each group of variables $x_1^a, x_2^a, \dots, x_{n_a}^a$. The numbers w_i are determined by the annihilation conditions satisfied by the vector $\mathbf{y}$.

In other words, we have the decomposition

$$(4.7) \quad J_{\mathbf{y}}^* = \bigoplus_{\mathbf{x} \in \Gamma(A)} J_{\mathbf{y}}^*(\mathbf{x})$$

and $J_{\mathbf{y}}^*(\mathbf{x})$ can be identified with the space of correlation functions of degree $\mathbf{x}$. Using Proposition 4.1 we obtain the following formula of Gordon type.

Proposition 4.2. *The character of the space $J_{\mathbf{y}}$ is given by the formula*

$$(4.8) \quad \text{ch } J_{\mathbf{y}}(q, \mathbf{z}) = \sum_{\mathbf{n} \in \mathbb{Z}^\ell} \mathbf{z}^{\mathbf{n}} q^{\frac{1}{2}\mathbf{n}A \cdot \mathbf{n} + \mathbf{v} \cdot \mathbf{n}} \prod_{a \in I} \frac{1}{(q)_{\mathbf{e}_a \cdot \mathbf{n}}}$$

where $\mathbf{v} = -\frac{1}{2}\text{diag}(A) + (w_1, w_2, \dots, w_\ell)$ and $\mathbf{z}^{\mathbf{n}} = z_1^{n_1} z_2^{n_2} \dots z_\ell^{n_\ell}$.

Let $\mathbf{N}$ be an ℓ -dimensional vector with integer components. Define the subspaces $W^A(\mathbf{N}) \subset \overline{\mathfrak{B}}(A)$ as follows:

$$(4.9) \quad W^A(\mathbf{N}) \text{ is spanned by } \{V(\mathbf{e}_a)_n \mid n \geq N_a, a \in I\}.$$

Then $W^A(\mathbf{N})$ -coinvariants of $J_{\mathbf{y}}^*$ can be easily calculated in terms of correlation functions. Namely the dual space consists of functions given in Proposition 4.1 with a restriction on the degrees in each variable. This gives the following

Proposition 4.3. *The character of $J_{\mathbf{y}}[W^A(\mathbf{N})] = J_{\mathbf{y}}^*/W^A(\mathbf{N})J_{\mathbf{y}}^*$ is given by the formula*

$$(4.10) \quad \text{ch } J_{\mathbf{y}}[W^A(\mathbf{N})](q, \mathbf{z}) = \sum_{\mathbf{n} \in \mathbb{Z}^\ell} \mathbf{z}^{\mathbf{n}} q^{\frac{1}{2}\mathbf{n}A \cdot \mathbf{n} + \mathbf{v} \cdot \mathbf{n}} \prod_{a \in I} \left[\begin{matrix} \mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n}A + \mathbf{w}) \\ \mathbf{e}_a \cdot \mathbf{n} \end{matrix} \right]_q$$

where $\mathbf{w} = (w_1, w_2, \dots, w_\ell)$.

4.2. Consider the family of algebras $\overline{\mathfrak{A}}_p^d$ obtained from the algebra $\mathfrak{A}_p^{\text{ass}}$ by the following recursive procedure.

Introduce a filtration on $\mathfrak{A}_p^{\text{ass}}$ by assigning degree 1 to the generators $\mathcal{G}^+$ and $\mathcal{G}^-$. The associated graded algebra $\overline{\mathfrak{A}}_p^0$ is generated by $\overline{\mathcal{G}}^+, \overline{\mathcal{G}}^-$. Define the currents $\overline{\mathcal{H}}^m$ by the following OPE:

$$\overline{\mathcal{G}}^+(z)\overline{\mathcal{G}}^-(w) = \frac{\overline{\mathcal{H}}^0}{(z-w)^{p-1}} + \dots + \frac{\overline{\mathcal{H}}^{p-2}}{z-w} + \text{regular terms}.$$

Note that the original algebra $\mathfrak{A}_p^{\text{ass}}$ is the central extension of $\overline{\mathfrak{A}}_p^0$.

At the next step we assign degrees 1 to $\overline{\mathcal{G}}^+, \overline{\mathcal{G}}^-$ and $\overline{\mathcal{H}}^0$. After taking graded object we obtain the algebra $\overline{\mathfrak{A}}_p^1$ whose generators we denote abusing notations again by $\overline{\mathcal{G}}^+, \overline{\mathcal{G}}^-$ and $\overline{\mathcal{H}}^0$. The algebra $\overline{\mathfrak{A}}_p^1$ is generated by currents $\overline{\mathcal{G}}^+, \overline{\mathcal{G}}^-$ and $\overline{\mathcal{H}}^0$ that have OPEs with the following properties

$$(4.11) \quad \overline{\mathcal{G}}^\pm(z)\overline{\mathcal{G}}^\pm(w) \sim (z-w)^p, \quad \overline{\mathcal{G}}^\pm(z)\overline{\mathcal{H}}^0(w) \sim (z-w)^1, \quad \overline{\mathcal{H}}^0(z)\overline{\mathcal{H}}^0(w) \sim (z-w)^2$$

$$(4.12) \quad \overline{\mathcal{G}}^+(z)\overline{\mathcal{G}}^-(w) = \frac{\overline{\mathcal{H}}^1}{(z-w)^{p-2}} + \dots + \frac{\overline{\mathcal{H}}^{p-2}}{z-w} + \text{regular terms}$$

To make the step $\overline{\mathfrak{A}}_p^{d-1} \rightarrow \overline{\mathfrak{A}}_p^d$ we introduce a filtration by attaching degree 1 to $\overline{\mathcal{G}}^+$, $\overline{\mathcal{G}}^-$ and $\overline{\mathcal{H}}^0$, $\dots$, $\overline{\mathcal{H}}^{d-1}$. As the result we obtain the algebra $\overline{\mathfrak{A}}_p^d$ with generators $\overline{\mathcal{G}}^+$, $\overline{\mathcal{G}}^-$ and $\overline{\mathcal{H}}^i$, $0 \leq i \leq d-1$ that have the following OPEs

(4.13)

$$\overline{\mathcal{G}}^\pm(z)\overline{\mathcal{G}}^\pm(w) \sim (z-w)^p, \quad \overline{\mathcal{G}}^\pm(z)\overline{\mathcal{H}}^m(w) \sim (z-w)^{\mathcal{A}_{\pm m}}, \quad \overline{\mathcal{H}}^n(z)\overline{\mathcal{H}}^m(w) \sim (z-w)^{\mathcal{A}_{nm}}$$

(4.14)

$$\overline{\mathcal{G}}^+(z)\overline{\mathcal{G}}^-(w) = \frac{\overline{\mathcal{H}}^d}{(z-w)^{p-d-1}} + \dots + \frac{\overline{\mathcal{H}}^{p-2}}{z-w} + \text{regular terms}$$

determined by the following $(d+2) \times (d+2)$ matrix

$$\mathcal{A} = \begin{pmatrix} p & -p+d+1 & 1 & 2 & 3 & \dots & d \\ -p+d+1 & p & 1 & 2 & 3 & \dots & d \\ 1 & 1 & 2 & 2 & 2 & \dots & 2 \\ 2 & 2 & 2 & 4 & 4 & \dots & 4 \\ 3 & 3 & 2 & 4 & 6 & \dots & 6 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ d & d & 2 & 4 & 6 & \dots & 2d \end{pmatrix}$$

where the first row and column corresponds to the index $+$ the second ones to $-$ the third to 0 and so on till $d-1$.

Proposition 4.4. *There is the surjective homomorphism*

$$\overline{\mathfrak{B}}(\mathcal{A}) \rightarrow \overline{\mathfrak{A}}_p^d$$

acting on generators as

$$V(\mathbf{e}_+, z) \rightarrow \overline{\mathcal{G}}^+(z), \quad V(\mathbf{e}_-, z) \rightarrow \overline{\mathcal{G}}^-(z), \quad V(\mathbf{e}_i, z) \rightarrow \overline{\mathcal{H}}^i(z), \quad 0 \leq i \leq d-1.$$

Proof. From (4.13) we can deduce that all relations between $V(\mathbf{e}_a, z)$ hold for their images in $\overline{\mathfrak{A}}_p^d$. So (4.16) defines a homomorphism. As the image contains all the generators of $\overline{\mathfrak{A}}_p^d$, this map is surjective. $\square$

Note that the map (4.16) is graded. Namely, it preserves the grading with respect to $\mathcal{L}_0$ and specialize the Γ -grading of $\overline{\mathfrak{B}}(\mathcal{A})$ into $\mathcal{H}^0$ -grading of $\overline{\mathfrak{A}}_p^d$. This specialization is defined by the functional $\mathbf{u}$ on Γ such that $\mathbf{u}(\mathbf{e}_+) = 1$, $\mathbf{u}(\mathbf{e}_-) = -1$, $\mathbf{u}(\mathbf{e}_i) = 0$ (see (2.21)).

Any representation $\mathbf{l}_r$ of $\overline{\mathfrak{A}}_p^{\text{ass}}$ contains the highest weight vector. Therefore the filtration in $\overline{\mathfrak{A}}_p^{\text{ass}}$ determines a filtration in $\mathbf{l}_r$ and associated graded algebra $\overline{\mathfrak{A}}_p^0$ acts in the associated graded module $\bar{\mathbf{l}}_r^0$. We can repeat this procedure with $\bar{\mathbf{l}}_r^0$ and the image of the highest weight vector. In this way we obtain the family of representations $\bar{\mathbf{l}}_r^d$ of algebras $\overline{\mathfrak{A}}_p^d$. Evidently characters of all $\bar{\mathbf{l}}_r^d$, $0 \leq d \leq p-2$ are equal to each other and equal to character of $\mathbf{l}_r$ given by (2.16).

Note that (4.16) allows us to treat any $\overline{\mathfrak{A}}_p^d$ -module as a $\overline{\mathfrak{B}}(\mathcal{A})$ -module. According to this we have

Proposition 4.5. *Let $\mathbf{y} = r\mathbf{u}$. Then there is the surjective mapping of $\overline{\mathfrak{B}}(\mathcal{A})$ -modules*

$$\mathbf{J}_{\mathbf{y}} \rightarrow \bar{\mathbf{l}}_r^d$$

mapping the vector $|\mathbf{y}\rangle$ into the highest weight vector.

Proof. It follows from the observation that the highest weight vector of $\bar{\mathbf{l}}_r^d$ satisfy the same annihilation conditions as $|\mathbf{y}\rangle$. $\square$

We will show (see Corollary 6.2) that (4.18) is indeed an isomorphism. Note that this map is bi-graded with respect to $\mathcal{L}_0$ and $\mathbf{u}(\Gamma)$.

Define the subspaces $W^d(\mathbf{N}) \subset \bar{\mathfrak{A}}_p^d$ recursively from $W(\mathbf{N})$ as the adjoint graded subspaces.

Proposition 4.6. *Let $0 \leq d \leq p-1$. Then $W^d(\mathbf{N})$ contains the image of $W^{\mathcal{A}}(\mathbf{N})$ under the map (4.16).*

Proof. Note that $W(\mathbf{N})$ contains linear combinations of the form

$$\mathcal{H}_s^i + c_1 \mathcal{H}_{s-1}^{i-1} + \cdots + c_i \mathcal{H}_s^0 + c_{i+1}$$

for $s \geq N_i - i$. Therefore $W^d(\mathbf{N})$ contains $\bar{\mathcal{H}}_s^i$ for $s \geq N_i - i$ hence (according to (2.8) and (4.2)) it contains the image of $W^{\mathcal{A}}(\mathbf{N})$. $\square$

The most important case of this statement is $d = p-1$, where the condition on $\mathbf{N}$ is empty.

Corollary 4.7. *Under the conditions of Proposition 4.6 we have the surjective map*

$$(4.19) \quad J_{\mathbf{ru}}[W^{\mathcal{A}}(\mathbf{N})] \rightarrow \bar{\mathbf{l}}_r^d[W(\mathbf{N})]$$

This map is an isomorphism only under some additional conditions (see Corollary 6.2).

The deformation argument (see for example [FKLMM2]) implies that for any filtered representation V of filtered (sub)algebra W we have

$$\dim(\text{gr } V[\text{gr } W]) \geq \dim V[W].$$

In our case it implies

$$(4.20) \quad \dim \bar{\mathbf{l}}_r^d[W^d(\mathbf{N})] \geq \dim \bar{\mathbf{l}}_r^{d-1}[W^{d-1}(\mathbf{N})] \geq \cdots \geq \dim \bar{\mathbf{l}}_r^0[W^0(\mathbf{N})] \geq \dim \mathbf{l}_r[W(\mathbf{N})]$$

Taking (4.19), (4.20) and Proposition 3.4 together we obtain the *main inequality*

Proposition 4.8. *For $0 \leq d \leq p-1$ we have*

$$(4.21) \quad \dim J_{\mathbf{ru}}[W^{\mathcal{A}}(\mathbf{N})] \geq \dim \mathbf{l}_r[W(\mathbf{N}^{i^1, j^1}, \mathbf{N}^{i^2, j^2}, \dots, \mathbf{N}^{i^k, j^k}; 0)] \geq \\ \dim \mathbf{l}_r[W(\mathbf{N}^{i^1, j^1}, \mathbf{N}^{i^2, j^2}, \dots, \mathbf{N}^{i^k, j^k}; \mathbf{a})] \geq d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k]$$

5. COMBINATORICS OF THE CHARACTERS

5.1. Following [SW] introduce q -supernomial coefficients as follows

$$(5.1) \quad \binom{\mathbf{L}}{a}_q^{k+1} = \sum_{\substack{n_1, \dots, n_k \in \mathbb{Z} \\ n_1 + \dots + n_k = a}} q^{\sum_{i=2}^k n_{i-1}(-n_i + \sum_{j=i}^k L_j)} \begin{bmatrix} L_k \\ n_k \end{bmatrix}_q \begin{bmatrix} L_{k-1} + n_k \\ n_{k-1} \end{bmatrix}_q \cdots \begin{bmatrix} L_1 + n_2 \\ n_1 \end{bmatrix}_q,$$

where $\mathbf{L} = (L_1, L_2, \dots, L_k)$ is a k -dimensional vector with nonnegative integer entries. The sum in the RHS is a well-defined polynomial in q because q -binomial coefficients vanish whenever the bottom index becomes greater than the top one.

Let $\mathbf{N} = (N_+, N_-; N_0, \dots, N_{d-1})$ be a vector in $\mathbb{Z}^{d+2}$; let $\mathbf{N}' = (N_0, \dots, N_{d-1}, N_+ + N_-)$ and let $\mathbf{L} = \mathbf{N}' T_{d+1}^{-1}$. The aim of this section is to prove the following polynomial identity.

Theorem 5.1. *Let $0 \leq d < 2p - 2$. Suppose that $L_i \geq 0$ and additionally*

$$(5.2) \quad 2N_+ - N_{d-1} \geq -(2p - d - 2), \quad 2N_- - N_{d-1} \geq -(2p - d - 2).$$

Then

$$(5.3) \quad \sum_{\mathbf{n} \in \mathbb{Z}^\ell} z^{n_+ - n_-} q^{\frac{1}{2} \mathbf{n} \mathbf{A} \cdot \mathbf{n}} \prod_{a \in I} \left[\frac{\mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n} \mathbf{A})}{\mathbf{e}_a \cdot \mathbf{n}} \right]_q = \sum_{a \in \mathbb{Z}} z^a q^{pa^2/2} \binom{\mathbf{L}}{pa + N_-}_q^{d+2}.$$

where $I = \{+, -, 0, 1, \dots, d-1\}$

We call (5.2) well-balanced conditions. Without them identity (5.3) can fail.

To eliminate the well-balanced conditions we use an improved version of q -binomial coefficients:

$$(5.4) \quad \begin{bmatrix} n \\ m \end{bmatrix}_q^+ = \begin{cases} \begin{bmatrix} n \\ m \end{bmatrix}_q & n \geq 0 \\ (-1)^{n-m} q^{-\frac{(n-m)^2 + (n-m)}{2}} \begin{bmatrix} -m-1 \\ -n-1 \end{bmatrix}_{q^{-1}} & n < 0 \end{cases}$$

In particular, for $q \rightarrow 1$ the integer $\begin{bmatrix} n \\ m \end{bmatrix}_1^+$ is the coefficient before z^{-m} in the Laurent expansion of $(1 + z^{-1})^n$ at $z = 0$.

This definition is motivated by the following relations.

Proposition 5.2. *The polynomials $\begin{bmatrix} n \\ m \end{bmatrix}_q^+$ are uniquely defined by the q -Pascal identities*

$$(5.5) \quad \begin{bmatrix} n \\ m \end{bmatrix}_q^+ = q^m \begin{bmatrix} n-1 \\ m \end{bmatrix}_q^+ + \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}_q^+ \quad \text{for any } n, m$$

$$(5.6) \quad \begin{bmatrix} n \\ m \end{bmatrix}_q^+ = \begin{bmatrix} n-1 \\ m \end{bmatrix}_q^+ + q^{n-m} \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}_q^+ \quad \text{for any } n, m$$


and conditions

$$(5.7) \quad \begin{bmatrix} m \\ m \end{bmatrix}_q^+ = 1, \quad \begin{bmatrix} n \\ 0 \end{bmatrix}_q^+ = 0 \quad \text{for } n < 0$$

which need to be checked only for certain m and n .

Proof. Indeed any set of polynomials $\begin{bmatrix} n \\ m \end{bmatrix}_q^?$ satisfying (5.5) and (5.6) can be expressed in the following way:

$$\begin{bmatrix} n \\ m \end{bmatrix}_q^? = \begin{cases} P_1 \cdot (-1)^{n-m} q^{-\frac{(n-m)^2 + (n-m)}{2}} \begin{bmatrix} -m-1 \\ n-m \end{bmatrix}_{q^{-1}} & n < 0, m < 0 \\ P_2 \cdot (-1)^m q^{-\frac{m^2 + m}{2}} \begin{bmatrix} -(n-m)-1 \\ m \end{bmatrix}_{q^{-1}} & n < 0, m \geq 0 \\ (P_1 + P_2) \cdot \begin{bmatrix} n \\ m \end{bmatrix}_q & n \geq 0 \end{cases}$$



where P_1 and P_2 are certain polynomials in q .

In our case we need only to check that $P_1 = 1$ and $P_2 = 0$.

□

Note that for usual q -binomial coefficients (5.5) and (5.6) fails for $n = m = 0$.

Theorem 5.3. *Let $0 \leq d < 2p - 2$. Suppose that $L_i \geq 0$. Then*

$$(5.8) \quad \sum_{\mathbf{n} \in \mathbb{Z}^\ell} z^{n_+ - n_-} q^{\frac{1}{2}\mathbf{n}\mathbf{A} \cdot \mathbf{n}} \prod_{a \in I} \left[\begin{matrix} \mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n}\mathbf{A}) \\ \mathbf{e}_a \cdot \mathbf{n} \end{matrix} \right]_q^+ = \sum_{a \in \mathbb{Z}} z^a q^{pa^2/2} \left(\begin{matrix} \mathbf{L} \\ pa + N_- \end{matrix} \right)_q^{d+2}.$$

Emphasize that in the left hand side the sum goes over all integer vectors.

5.2. Here we show that the left hand side of (5.8) is a well defined polynomial and that theorem 5.3 implies theorem 5.1.

Let $P_{\mathbf{N}}(\mathbf{n}) = \prod_{a \in I} \left[\begin{matrix} \mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n}\mathbf{A}) \\ \mathbf{e}_a \cdot \mathbf{n} \end{matrix} \right]_q^+$. We need to prove that if $L_i \geq 0$ for any i then there is only finite number of vectors $\mathbf{n}$ such that $P_{\mathbf{N}}(\mathbf{n}) \neq 0$. And if additionally we impose the well-balance conditions (5.2) then $P_{\mathbf{N}}(\mathbf{n}) \neq 0$ implies $n_a \geq 0$, $a = +, -, 0, 1, \dots, d-1$, where $n_a = \mathbf{e}_a \cdot \mathbf{n}$.

Proposition 5.4. *If $P_{\mathbf{N}}(\mathbf{n}) \neq 0$ then*

$$(5.9) \quad N_a \geq \mathbf{e}_a \cdot \mathbf{n}\mathbf{A} \quad \text{for any } a = +, -, 0, 1, 2, \dots, d-1;$$

$$(5.10) \quad N_a < \mathbf{e}_a \cdot (\mathbf{n}\mathbf{A} - \mathbf{n}) \quad \text{if } n_a < 0.$$

Proof. Condition (5.9) follows from the fact that $\left[\begin{smallmatrix} n \\ m \end{smallmatrix} \right]_q^+ = 0$ if $n < m$. Condition (5.10) follows from the fact that $\left[\begin{smallmatrix} n \\ m \end{smallmatrix} \right]_q^+ = 0$ if $m < 0$ and $n \geq 0$. $\square$

Lemma 5.5. *Let $1 \leq i \leq d$. If $L_i \geq 0$ and $n_{i-1} < 0$ then $P_{\mathbf{N}}(\mathbf{n}) = 0$.*

Proof. Suppose that $n_{i-1} < 0$ and $P_{\mathbf{N}}(\mathbf{n}) \neq 0$. Then adding (5.10) for $a = i-1$ twice and subtracting (5.9) for $a = i-2$ (if $i > 1$) and $a = i$ (if $i < d$; if $i = d$ then for $a = +$ and $a = -$) we obtain $L_i < 0$. $\square$

Lemma 5.6. *If $L_{d+1} \geq 0$ and $n_+ < 0$, $n_- < 0$ then $P_{\mathbf{N}}(\mathbf{n}) = 0$.*

Proof. Proof is similar. Here we add (5.10) for $a = +$ and for $a = -$ then subtract (5.9) for $a = d-1$. $\square$

Lemma 5.7. *Suppose that $n_+ < 0$ and $n_- \geq 0$. Then $P_{\mathbf{N}}(\mathbf{n}) = 0$ for*

$$n_+ \leq C_+ = (2N_+ - N_{d-1})/(2p - d - 2).$$

Proof. Suppose that $n_+ < 0$ and $P_{\mathbf{N}}(\mathbf{n}) \neq 0$. Then adding (5.10) for $a = +$ twice and subtracting (5.9) for $a = d-1$ we obtain

$$2N_+ - N_{d-1} < (2p - d - 2)n_+ - (2p - d - 2)n_-.$$

Taking into account $n_- \geq 0$ and $2p - d - 2 > 0$ we deduce $(2p - d - 2)n_+ > 2N_+ - N_{d-1}$. $\square$

Clearly, we have the similar statement for $n_- < 0$ and $n_+ \geq 0$.

So if $L_i \geq 0$ for any i and the well-balanced conditions are satisfied then $P_{\mathbf{N}}(\mathbf{n}) \neq 0$ only if $n_a \geq 0$ for any $a = +, -, 0, \dots, d-1$, hence theorem 5.3 implies theorem 5.1.

Under the conditions of theorem 5.3 we have $n_i \geq 0$ for $i = 0 \dots d-1$ and $n_+ > C_+$, $n_- > C_-$ otherwise $P_{\mathbf{N}}(\mathbf{n}) = 0$. Adding (5.9) for $a = +$ and $a = -$, we obtain

$$N_+ + N_- \geq (d+1)n_+ + (d+1)n_- + 2n_0 + 4n_1 + 6n_2 + \dots$$

As all the coefficient in the right hand side of this inequality are positive, there exists only finite number of such vectors $\mathbf{n}$.

5.3. Here we compare the left hand side $\chi_{p,d}^L[\mathbf{N}](q, z)$ and the right hand side $\chi_{p,d}^R[\mathbf{L}, N_-](q, z)$ of (5.8).

Note that for q -supernomial coefficients there is an identity

$$\binom{\mathbf{L} + \mathbf{e}_k}{a}_q^{k+1} = \binom{\mathbf{L} + \mathbf{e}_{k-1}}{a-1}_q^{k+1} + q^a \binom{\mathbf{L}}{a}_q^k.$$

Therefore, for the right hand side we have

$$(5.11) \quad \chi_{p,d}^R[\mathbf{L} + \mathbf{e}_{d+1}, N_-] = \chi_{p,d}^R[\mathbf{L} + \mathbf{e}_d, N_- - 1] + z^{-1} q^{N_- - p/2} \chi_{p,d}^R[\mathbf{L}, N_- - p].$$

Also as $\binom{(L_1, \dots, L_{k-1}, 0)}{a}_q^{k+1} = \binom{(L_1, \dots, L_{k-1})}{a}_q^k$, we have

$$(5.12) \quad \chi_{p,d}^R[(L_1, \dots, L_d, 0), N_-] = \chi_{p,d-1}^R[(L_1, \dots, L_d), N_-].$$

The idea of the proof is to check (5.11) and (5.12) for the left hand side.

Proposition 5.8. *Let A be a symmetric $m \times m$ matrix with integer components, let $\mathbf{u}, \mathbf{N}$ be vectors in $\mathbb{Z}^m$. Consider the polynomial*

$$(5.13) \quad \chi_{A,\mathbf{u}}[\mathbf{N}](q, z) = \sum_{\mathbf{n} \in \mathbb{Z}^m} z^{\mathbf{u} \cdot \mathbf{n}} q^{\frac{1}{2} \mathbf{n} A \cdot \mathbf{n}} \prod_a \left[\frac{\mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n} A)}{\mathbf{e}_a \cdot \mathbf{n}} \right]_q^+.$$

Then for any a we have

$$(5.14) \quad \chi_{A,\mathbf{u}}[\mathbf{N}] = \chi_{A,\mathbf{u}}[\mathbf{N} - \mathbf{e}_a] + z^{u_a} q^{N_a - A_{aa}/2} \chi_{A,\mathbf{u},\mathbf{v},\mathbf{w}}[\mathbf{N} - \mathbf{e}_a A].$$

Proof. Identity (5.14) follows from the identity (5.6) applied to the factor $\left[\frac{\mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n} A)}{\mathbf{e}_a \cdot \mathbf{n}} \right]_q^+$ in each summand. $\square$

Applying (5.14) with $a = -$ to $\chi_{p,d}^L[\mathbf{N}]$, we check (5.11) for the left hand side. Note that for $a = 0, \dots, d-1$ (5.14) implies recurrence relations for the q -supernomial coefficients same as in [SW].

To check (5.11) for the left hand side we need the following identity.

Lemma 5.9. *There is the following relation*

$$(5.15) \quad \begin{bmatrix} N \\ n \end{bmatrix}_q^+ \begin{bmatrix} M \\ m \end{bmatrix}_q^+ = \sum_{l \in \mathbb{Z}} q^{(n-l)(m-l)} \begin{bmatrix} N-m \\ n-l \end{bmatrix}_q^+ \begin{bmatrix} M-n \\ m-l \end{bmatrix}_q^+ \begin{bmatrix} N+M-n-m+l \\ l \end{bmatrix}_q^+$$

Proof. First let us check that the sum in the right hand side is indeed finite.

Note that if $N + M - n - m < 0$ then the third factor is always zero, so entire sum is zero. If $N + M - n - m \geq 0$ then either $N - m \geq 0$ or $M - n \geq 0$. In the first case we have non-zero summands only if $n + m - N \leq l \leq n$, in the second case only if $n + m - M \leq l \leq m$.

Let us denote the polynomial in the right hand side by $\text{RHS}_{n,m}^{N,M}$. The idea of proof is to apply Proposition 5.2 to $\text{RHS}_{n,m}^{N,M}$ as a set of polynomials indexed by N and n . It means that we prove that

$$(5.16) \quad \text{RHS}_{n,m}^{N,M} = q^n \text{RHS}_{n,m}^{N-1,M} + \text{RHS}_{n-1,m}^{N-1,M}, \quad \text{RHS}_{n,m}^{N,M} = \text{RHS}_{n,m}^{N-1,M} + q^{N-n} \text{RHS}_{n-1,m}^{N-1,M}$$

for any N, n, M, m and

$$(5.17) \quad \text{RHS}_{n,m}^{n,M} = \begin{bmatrix} M \\ m \end{bmatrix}_q^+ \quad \text{RHS}_{0,m}^{N,M} = 0$$

for any M, m and certain $N < 0, n$.

To prove (5.16) fix for a moment N, M, n, m and introduce the notation

$$\begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} = q^{(n-l)(m-l)} \begin{bmatrix} N-m+a \\ n-l+d \end{bmatrix}_q^+ \begin{bmatrix} M-n+b \\ m-l+e \end{bmatrix}_q^+ \begin{bmatrix} N+M-n-m+l+c \\ l+f \end{bmatrix}_q^+.$$

Then substitution $l+1$ into l leads to the identity

$$\sum_{l \in \mathbb{Z}} \begin{bmatrix} a & b & c-1 \\ d+1 & e+1 & f-1 \end{bmatrix} = \sum_{l \in \mathbb{Z}} q^{2l-m-n+1} \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}.$$

Using this identity together with (5.5) we obtain

$$\begin{aligned} \text{RHS}_{n,m}^{N,M} &= \sum \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \sum q^{n-l} \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} + \sum \begin{bmatrix} -1 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} = \\ &= \sum q^n \begin{bmatrix} -1 & 0 & -1 \\ 0 & 0 & 0 \end{bmatrix} + \sum q^{n-l} \begin{bmatrix} -1 & 0 & -1 \\ 0 & 0 & -1 \end{bmatrix} + \sum \begin{bmatrix} -1 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} = \\ &= \sum q^n \begin{bmatrix} -1 & 0 & -1 \\ 0 & 0 & 0 \end{bmatrix} + \sum q^{l-m} \begin{bmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \end{bmatrix} + \sum \begin{bmatrix} -1 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} = \\ &= \sum q^n \begin{bmatrix} -1 & 0 & -1 \\ 0 & 0 & 0 \end{bmatrix} + \sum q^{l-m} \begin{bmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \end{bmatrix} = q^n \text{RHS}_{n,m}^{N-1,M} + \text{RHS}_{n-1,m}^{N-1,M}, \end{aligned}$$

where sums go over $l \in \mathbb{Z}$. Proof of the second part of (5.16) is similar.

To check (5.17) we take $n = M$ and any negative $N < m - M$. We have only one non-zero summand in $\text{RHS}_{M,m}^{M,M}$, namely for $l = m$, and it coincides with the q -binomial given in (5.17). If $N < m - M$ and $n = 0$ then $N + M - m - n < 0$ and, as described above, the result is zero.

So proposition (5.2) implies the lemma. $\square$

Applying (5.15) to the factors in $\chi_{p,d-1}^L[(N_+, N_-; N_0 \dots, N_{d-2})]$ corresponding to n_+ and n_- we obtain

$$\chi_{p,d-1}^L[(N_+, N_-; N_0 \dots, N_{d-2})] = \chi_{p,d}^L[(N_+, N_-; N_0 \dots, N_{d-2}, N_+ + N_-)],$$

that is, (5.12) for the left hand side.

Now we can prove theorem 5.3 by induction on d and L_{d+1} . To complete the proof we need only to check the case $d = 0$ manually.

Lemma 5.10. *Suppose that $M + S \geq 0$. Then*

$$(5.18) \quad \sum_{k \in \mathbb{Z}} q^{k^2 + ak} \begin{bmatrix} M \\ a + k \end{bmatrix}_q^+ \begin{bmatrix} S \\ k \end{bmatrix}_q^+ = \begin{bmatrix} M + S \\ S + a \end{bmatrix}_q$$

Proof. First let us check that the sum in the left hand side is indeed finite. As $M + S \geq 0$ we have either $M \geq 0$ or $S \geq 0$. In the first case we have non-zero summands only if $-a \leq k \leq M - a$, in the second case only if $0 \leq k \leq S$.

Let us denote the polynomial in the left hand side by $\text{LHS}_a^{M,S}$. Clearly, we have $\text{LHS}_a^{M,S} = \text{LHS}_{-a}^{S,M}$ and the same identity for the right hand side. Therefore we can restrict ourselves to the case $S \geq 0$ and prove that

$$(5.19) \quad \text{LHS}_a^{M,S} = \begin{bmatrix} M + S \\ S + a \end{bmatrix}_q^+ \quad \text{for } S \geq 0 \text{ and any } M.$$

Applying (5.6) to the second factor in $\text{LHS}_a^{M,S}$, we obtain

$$\text{LHS}_a^{M,S} = \text{LHS}_a^{M,S-1} + q^{S+a} \text{LHS}_{a+1}^{M,S-1},$$

that is (5.5) for the right hand side of (5.19). Therefore it remains to check (5.19) for $S = 0$.

If $S = 0$ then the only non-zero summand in $\text{LHS}_a^{M,S}$ is for $k = 0$ and it is equal to the right hand side of (5.19) $\square$

The case $d = 0$ of Theorem 5.3 can be obtained by substituting $M = N_+ - (p-1)(n_+ - n_-)$, $S = N_- + (p-1)(n_+ - n_-)$, $a = n_+ - n_-$ and $k = n_-$ in (5.18).

6. DIMENSION AND CHARACTER FORMULAS FOR THE COINVARIANTS

Theorem 6.1. *For any $\mathbf{a} \in \mathbb{C}^k$ we have*

$$(6.1) \quad \dim \mathbb{I}_r[W(\mathbf{N}^{i^1, j^1}, \mathbf{N}^{i^2, j^2}, \dots, \mathbf{N}^{i^k, j^k}; \mathbf{a})] = d_{p,r}[i^1, j^1; i^2, j^2; \dots; i^k, j^k].$$

Proof. First, note that due to the action of automorphisms $\mathbf{U}_\theta$ we can consider only coinvariants of the representation $\mathbb{I}_0$.

The main idea of the proof is to check that the main inequality (4.21) is indeed an equality. Here we have a chain of inequalities, where we know the first element by Proposition 4.3 and the last element. As soon as we check that these two numbers are equal we obtain that other numbers in (4.21) are also equal.

Suppose that the vector $\mathbf{N}$ satisfies the well-balance conditions (5.2). Then this equality follows from theorem 5.1 by setting $z = 1$ and $q = 1$. But if conditions (5.2) are not satisfied then this equality can be violated. So we should deduce general case from the well-balanced one.

To this end consider the family of automorphisms $U_{p\theta}$. They preserve the representation l_0 and permute the subalgebras $W(\mathbf{N})$. Namely, $U_{p\theta}$ preserves $N_0, \dots, N_{d-1}$, increases N_+ by $p\theta$ and decreases N_- by $p\theta$.

Recall that $L_p = N_+ + N_- - N_{p-2}$.

First consider the case $L_p > 0$. Then we should set $d = p - 1$. Let $\theta \in \mathbb{Z}$ be the minimal number such that $2N_+ + p\theta - N_{p-2} \geq -p + 1$. So we have $2N_+ + p\theta - N_{p-2} \leq p$ and as $L_p \geq 1$ we obtain $2N_- - p\theta - N_{p-2} > -p + 1$. Hence after applying the automorphism $U_{p\theta}$ the well-balance conditions will be satisfied.

In the case $L_p = 0$ we set $d = p - 2$. Let as above $\theta \in \mathbb{Z}$ be the minimal number such that $2N_+ + p\theta - N_{p-3} \geq -p$. So we have $2N_+ + p\theta - N_{p-3} < p$ and as $L_{p-1} \geq 0$ we obtain $2N_- - p\theta - N_{p-3} > -p$. Hence after applying the automorphism $U_{p\theta}$ the well-balance conditions will be satisfied. $\square$

Corollary 6.2. *Let $0 \leq d \leq p - 1$. Let $\mathcal{A}$ be the matrix given by (4.15). Then we have*

$$(6.2) \quad J_{\mathbf{ru}} \cong \bar{l}_r^d.$$

If $N_d = N_{d+1} = \dots = N_{p-2} = N_+ + N_-$ and $\mathbf{N}$ satisfies the well-balance condition

$$2N_+ - N_{d-1} - 2r \geq -(2p - d - 2), \quad 2N_- - N_{d-1} + 2r \geq -(2p - d - 2)$$

then we have

$$(6.3) \quad J_{\mathbf{ru}}[W^{\mathcal{A}}(\mathbf{N})] \cong \bar{l}_r^d[W(\mathbf{N})]$$

So for the character of l_r we have a set of formulas indexed by $0 \leq d \leq p - 1$

$$\text{ch } l_r(q, z) = z^{-\frac{r}{p}} q^{\frac{r}{2p}(r-p+2)} \sum_{\mathbf{n} \in \mathbb{Z}^\ell} z^{\mathbf{u} \cdot \mathbf{n}} q^{\frac{1}{2} \mathbf{n} \cdot \mathcal{A} \cdot \mathbf{n} + \mathbf{v} \cdot \mathbf{n}} \prod_{a \in I} \frac{1}{(q)_{\mathbf{e}_a \cdot \mathbf{n}}},$$

where the set of indices is $I = \{+, -, 0, 1, 2, \dots, d-1\}$, $\mathbf{u} = (1, -1; 0, \dots, 0)$, $\mathbf{v} = (p/2 - r - 1)\mathbf{u}$.

Now discuss the character formulas for coinvariants.

Theorem 6.3. *Consider the character defined as $\text{ch } l_r[W(\mathbf{N})](q, z) = \text{tr } z^{\mathcal{H}_0} q^{\mathcal{L}_0}$.*

- (i) *Let the set of indices be $I = \{+, -, 0, 1, \dots, p-2\}$. Consider the vectors $\mathbf{u} = (1, -1; 0, \dots, 0)$, $\mathbf{v} = (p/2 - r - 1)\mathbf{u}$, $\mathbf{w} = r\mathbf{u}$. Then*

$$\text{ch } l_r[W(\mathbf{N})](q, z) = z^{-\frac{r}{p}} q^{\frac{r}{2p}(r-p+2)} \sum_{\mathbf{n} \in \mathbb{Z}^\ell} z^{\mathbf{u} \cdot \mathbf{n}} q^{\frac{1}{2} \mathbf{n} \cdot \mathcal{A} \cdot \mathbf{n} + \mathbf{v} \cdot \mathbf{n}} \prod_{a \in I} \left[\frac{\mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n} \mathcal{A} + \mathbf{w})}{\mathbf{e}_a \cdot \mathbf{n}} \right]_q^+,$$

where the notion $\left[\frac{n}{m} \right]_q^+$ is given by (5.4).

- (ii) *We have*

$$\text{ch } l_r[W(\mathbf{N})](q, z) = z^{-\frac{r}{p}} q^{\frac{r}{2p}(r-p+2)} \sum_{m \in \mathbb{Z}} z^m q^{\frac{p}{2}(m^2+m) - (r+1)m} \left(\frac{\mathbf{L}}{pm - r + N_-} \right)_q^{p+1}.$$

Proof. Note that U_θ acts on the character by substituting zq into z . Therefore it is enough to prove the theorem in the case $r = 0$.

First suppose that $\mathbf{N}$ satisfies the well-balance conditions (5.2). Then (i) follows from Corollary 6.2 and Proposition 4.3; the statement (ii) can be deduced from (i) by Theorem 5.1.

We know that general case can be reduced to the well-balanced case by the action of $U_{p\theta}$. As $U_{p\theta}$ acts on the RHS of (ii) by the same substitution zq^p into z , we have (ii) in general case. The statement (i) can be deduced from (ii) by Theorem 5.3. $\square$

Corollary 6.4. *Let $0 \leq d \leq p-2$. Suppose that $N_d = N_{d+1} = \dots = N_{p-2} = N_+ + N_-$. Let the set of indices be $I = \{+, -, 0, 1, \dots, d-1\}$ and, as above, $\mathbf{u} = (1, -1; 0, \dots, 0)$, $\mathbf{v} = (p/2 - r - 1)\mathbf{u}$, $\mathbf{w} = r\mathbf{u}$. Then*

$$\text{ch } l_r[W(\mathbf{N})](q, z) = z^{-\frac{r}{p}} q^{\frac{r}{2p}(r-p+2)} \sum_{\mathbf{n} \in \mathbb{Z}^\ell} z^{\mathbf{u} \cdot \mathbf{n}} q^{\frac{1}{2}\mathbf{n} \cdot \mathbf{A} \cdot \mathbf{n} + \mathbf{v} \cdot \mathbf{n}} \prod_{a \in I} \left[\begin{matrix} \mathbf{e}_a \cdot (\mathbf{N} + \mathbf{n} - \mathbf{n} \cdot \mathbf{A} + \mathbf{w}) \\ \mathbf{e}_a \cdot \mathbf{n} \end{matrix} \right]_q^+,$$

where the notion $\left[\begin{matrix} n \\ m \end{matrix} \right]_q^+$ is given by (5.4).

Acknowledgments. This work is supported by grants RFBR 00-15-96579, 01-01-00906, 01-02-16686 and 01-01-00546 and CRDF RP1-2254.

REFERENCES

- [D] C. Dong, *Vertex algebras associated with even lattice*, J. of Algebra 160 (1993).
- [FFr] B. Feigin, E. Frenkel, *Coinvariants of nilpotent subalgebras of the Virasoro algebra and partition identities*, Adv. Sov. Math. **16** (1993) p. 139–148, [hep-th/9301039](#).
- [FFu] B.L. Feigin, D.B. Fuchs, *Cohomology of some nilpotent subalgebras of the Virasoro and Kac-Moody algebras*, Geometry and Physics, Essays in honor of I.M. Gelfand on the occasion of his 75th birthday, eds. S. Gindikin and I.M. Singer, p. 209–235. North-Holland, 1991.
- [FKLMM1] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, *Combinatorics of the $\widehat{sl}_2$ spaces of coinvariants*, Transformation groups 1, 2001, [math-ph/9908003](#).
- [FKLMM2] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, *Combinatorics of the $\widehat{sl}_2$ spaces of coinvariants, II*, [math.QA/0009198](#)
- [FKLMM3] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, *Combinatorics of the $\widehat{sl}_2$ spaces of coinvariants, III*, [math.QA/0012190](#)
- [FL] B. Feigin and S. Loktev, *On generalized Kostka polynomials and the quantum Verlinde rule*, Differential topology, infinite-dimensional Lie algebras, and applications, Amer. Math. Soc. Transl. Ser. 2, 194 (1999), p. 61–79, [math.QA/9812093](#).
- [K] V. Kac, *Vertex Algebras for Beginners*, University Lecture Series, 10, 1997.
- [KKMM] R. Kedem, T. Klassen, B. McCoy, E. Melzer, *Fermionic sum representations for conformal field theory characters*. Phys. Lett. **B 307** (1993), p. 68–76.
- [SW] A. Schilling, S. Ole Warnaar, *Supernomial coefficients, polynomial identities and q -series*, The Ramanujan Journal 2 (1998) 459–494, [math.QA/9701007](#).
- [S] G. Segal, *Geometric aspect of quantum field theory*, Proceedings of the ICM at Kyoto, 1990.
- [V] E. Verlinde, *Fusion rules and modular transformations in 2D conformal field theory*, Nuclear Physics **B300** (1988) p. 360–376.

LANDAU INSTITUTE FOR THEORETICAL PHYSICS, RUSSIAN ACADEMY OF SCIENCES

INDEPENDENT UNIVERSITY OF MOSCOW

TAMM THEORY DIVISION, LEBEDEV PHYSICS INSTITUTE, RUSSIAN ACADEMY OF SCIENCES